

Honor fungicide

technology sheet

New multiple-mode-of-action fungicide controls key turf diseases while providing Intrinsic® plant health benefits.

- Up to 28 days of control for more than 20 turf diseases, including summer patch, anthracnose and dollar spot
- Multiple-mode-of-action control of dollar spot
- Unique Intrinsic plant health benefits¹ help mitigate environmental and mechanical stress²

Active ingredients

Pyraclostrobin – Group 11
Boscalid – Group 7

Formulation

Wettable granules

One case contains

2 x 1.42 kg jugs

Storage

Store in cool, dry, locked, ventilated area without a floor drain, away from food and animal feed.

¹ Intrinsic benefits refer to products that contain the active ingredient pyraclostrobin.

² All comparisons are to untreated, unless otherwise stated.

³ See label for more details on timing and restrictions.

⁴ Suppression only.

Crop

Turfgrass

Timing³

Apply preventatively, when conditions are favourable for disease.

Diseases controlled

Anthracnose (*Colletotrichum cereale*)
Bentgrass dead spot (*Ophiosphaerella agrostis*)
Brown patch (*Rhizoctonia solani, zea*)
Brown ring patch (*Rhizoctonia circinata* var. *circinata*)
Dollar spot (*Sclerotinia homoeocarpa*)
Fairy ring (Basidiomycete fungi)
Fusarium patch (*Microdochium nivale*)
Gray leaf spot (*Pyricularia grisea*)
Gray snow mold/Typhula blight (*Typhula incarnata, T. ishikariensis*)
Leaf spot (*Drechslera* spp., *Bipolaris* spp.)
Melting out (*Drechslera poae*)
Pink snow mold (*Microdochium nivale*)
Powdery mildew (*Erysiphe graminis*)
Pythium blight (*Pythium aphanidermatum*)
Pythium root dysfunction (*Pythium volutum, Pythium arrhenomanes, Pythium aristosporum*)
Rapid blight (*Labyrinthula terrestris*)
Red thread (*Laetisaria fuciformis*)
Rhizoctonia leaf or sheath spot (*R. oryzae, R. zea*)
Rust (*Puccinia* spp.)
Summer patch (*Magnaporthe poae*)
Take-all patch⁴ (*Gaeumannomyces graminis*)

Application rates⁴

One case of Honor™ fungicide treats 0.8 to 3 hectares (2 to 7 acres).

Crop	Application rate
Turfgrass	1.0 to 3.5 kg/ha (10.1 to 35 g/100 m ²) depending on disease

⁴ Refer to label for disease-specific rates and restrictions.

Mixing order

1. Ensure the spray tank is clean before use.
2. Fill the tank 1/2 full of water and start agitation.
3. Add the required amount of Honor fungicide to the tank while agitating.
4. If required, add the required amount of Premis® 200 F fungicide to tank.
5. Continue agitation while filling the remainder of the spray tank.
6. After use, clean the spray tank according to label precautions.

Application tips

Rainfastness – 1 hour.

Restricted entry interval – Once residues dry.

Resistance management – In order to reduce the risk of the development of fungicide resistance, adhere to the label instructions regarding the consecutive use of Honor fungicide or Group 7 or 11 fungicides that have a similar site of action on the same pathogens. Do not exceed the total number of sequential applications of Honor recommended. Do not exceed (4) four applications per season.

Additives

Do not use with organosilicate based adjuvants or turf grass injury may occur. Due to the large number of additives or adjuvants that may be used, neither the manufacturer nor the seller has determined whether Honor fungicide can be used safely with all additives.

Tank mixes

Fungicides: Premis 200 F

When used in a tank mix, follow all use directions and the most restrictive precautions on the individual product labels.

Contact **AgSolutions®** Customer Care or your BASF Sales Representative for additional information on supported tank mixes.


For more information

Call **AgSolutions** Customer Care at 1-877-371-BASF (2273)
or **better-turf.basf.ca**

Always read and follow label directions.


AgSolutions is a registered trade-mark of BASF Corporation; **HONOR** is a trade-mark and **INTRINSIC** is a registered trade-mark of BASF SE; **PREMIS** is a registered trade-mark of BASF Agro Trademarks GmbH; all used with permission by BASF Canada Inc. **HONOR** and **PREMIS** fungicides should be used in a preventative disease control program. © 2017 BASF Canada Inc.

Summer patch control



Source: BASF and 3rd party research

Anthracnose control



Source: BASF and 3rd party research, n=6