



We create chemistry

PRODUCT LABELS FOR

INSIGNIA DUO:

**INSIGNIA SC
FUNGICIDE**

+

**PREMIS 200 F
FUNGICIDE**

Container Label

GROUP	11	FUNGICIDE
-------	----	-----------

INSIGNIA® SC Fungicide

Broad spectrum fungicide for disease control on golf course turf

SUSPENSION CONCENTRATE

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENT: Pyraclostrobin 250 g/L

Contains 1,2-benzisothiazolin-3-one at 0.0038% and 2-methyl-4-isothiazolin-3-one at 0.0038%, as preservatives

OR

Contains 1,2-benzisothiazolin-3-one at 0.0075%, 2-methyl-4-isothiazolin-3-one at 0.0075%, and 2-bromo-2-nitropropane-1,3-diol at 0.024%, as preservatives

OR

Contains 1,2-benzisothiazolin-3-one at 0.0400%, 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0011%, 2-methyl-4-isothiazolin-3-one at 0.0004%, and 2-bromo-2-nitropropane-1,3-diol at 0.0240%, as preservatives

REGISTRATION NO. 32247

PEST CONTROL PRODUCTS ACT

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**



**DANGER POISON
SKIN IRRITANT**

READ THE LABEL AND BROCHURE BEFORE USING

NET CONTENTS: 0.1 L - Bulk

BASF Canada Inc.
5025 Creekbank Road
Building A, 2nd Floor
Mississauga, Ontario L4W 0B6
1-877-371-2273

INSIGNIA is a registered trade-mark of BASF Corporation, used under license by BASF Canada Inc.

PRECAUTIONS: Please refer to the booklet for the complete list of precautions.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

The patient should be treated symptomatically.

ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms, non-target terrestrial plants and small wild mammals. Observe buffer zones specified under DIRECTIONS FOR USE. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

STORAGE

1. Store in original tightly closed container.
2. Do not ship or store near food, feed, seed and fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.
4. Protect from freezing.
5. Store this product away from food or feed.

DISPOSAL

For Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for any further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

For Non-Returnable Containers

1. Triple or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial/territorial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial/territorial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

For Returnable-Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

INSIGNIA® SC Fungicide

Broad spectrum fungicide for disease control on golf course turf

SUSPENSION CONCENTRATE

COMMERCIAL (AGRICULTURAL)

ACTIVE INGREDIENT: Pyraclostrobin 250 g/L

Contains 1,2-benzisothiazolin-3-one at 0.0038% and 2-methyl-4-isothiazolin-3-one at 0.0038%, as preservatives

OR

Contains 1,2-benzisothiazolin-3-one at 0.0075%, 2-methyl-4-isothiazolin-3-one at 0.0075%, and 2-bromo-2-nitropropane-1,3-diol at 0.024%, as preservatives

OR

Contains 1,2-benzisothiazolin-3-one at 0.0400%, 5-chloro-2-methyl-4-isothiazolin-3-one at 0.0011%, 2-methyl-4-isothiazolin-3-one at 0.0004%, and 2-bromo-2-nitropropane-1,3-diol at 0.0240%, as preservatives

REGISTRATION NO. 32247

PEST CONTROL PRODUCTS ACT

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**



DANGER

POISON

SKIN IRRITANT

READ THE LABEL AND BROCHURE BEFORE USING

NET CONTENTS: 0.1 L - Bulk

BASF Canada Inc.
5025 Creekbank Road
Building A, 2nd Floor
Mississauga, Ontario L4W 0B6
1-877-371-2273

INSIGNIA is a registered trade-mark of BASF Corporation, used under license by BASF Canada Inc.

GENERAL INFORMATION

This package contains **INSIGNIA SC Fungicide**, a Suspension Concentrate (SC). The active ingredient in **INSIGNIA SC Fungicide** belongs to the class of fungicides, the strobilurins. Strobilurins are synthetic derivatives of a natural antifungal substance. **INSIGNIA SC Fungicide** has a mode of action that is effective against pathogens resistant to other fungicides groups.

INSIGNIA SC Fungicide inhibits spore germination and mycelial growth and sporulation of the fungus on the leaf surface. However, optimum disease control is achieved when **INSIGNIA SC Fungicide** is applied in a regularly scheduled protective spray program and is used in a rotation program with other fungicides.

DIRECTIONS FOR USE

INSIGNIA SC Fungicide is a broad-spectrum fungicide recommended for the control of many important diseases of turf grass. For maximum efficacy, **INSIGNIA SC Fungicide** should be applied preventively.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application

DO NOT apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE) medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply by air.

Spray Buffer Zones

Use of the following spray methods or equipment **DO NOT** require a spray buffer zone: hand-held or backpack sprayer and spot treatment.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs and wetlands) and estuarine/marine habitats.

Method of application	Pest	Spray Buffer zones (metres) required for the protection of:				
		Freshwater habitat of depths:		Estuarine/marine habitats of depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	Less than 1 m	Greater than 1 m	
Field sprayer*	Gray snow mold, typhula blight (low rate)	10	1	1	1	0
	Gray snow mold, typhula blight (high rate)	10	3	3	1	1
	Rust, gray leaf spot (low rate)	10	5	4	2	1
	Pink snow mold	15	2	2	1	0
	Leaf spot	15	5	4	2	1
	Anthrachnose, take-all patch, dollar spot (low rate)	15	5	4	2	1
	Pythium blight	20	5	5	3	1
	Brown patch, gray leaf spot, anthracnose, pythium root dysfunction, take-all patch, dollar spot (high rate)	20	10	5	3	1

*For field sprayer application, spray buffer zones can be reduced with the use of drift reducing spray shields. When using a spray boom fitted with a full shield (shroud, curtain) that extends to the crop canopy, the labelled buffer zone can be reduced by 70%. When using a spray boom where individual nozzles are fitted with cone-shaped shields that are no more than 30 cm above the crop canopy, the labelled spray buffer zone can be reduced by 30%.

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

All applications should be made according to the use directions indicated on Table 1. Failure to follow directions and precautions on this label may result in turf grass injury and/or inferior disease control.

INSIGNIA SC Fungicide is recommended for the control of Brown patch, Gray leaf spot, Gray snow mold, Leaf spot, Pink snow mold, Pythium blight, Pythium root dysfunction, Rust, Anthracnose, and for the suppression of Take-all patch and Dollar spot. For optimum control of Pink snow mold, tank mix **INSIGNIA SC Fungicide** with another snow mold fungicide (non-strobilurin). Apply **INSIGNIA SC Fungicide** in 40-120 litres of water per 1,000 square meters (400-1200 litres per hectare).

Rate: Use the application rates specified for each disease listed in Table 1.

Turf Grass Tolerance

Due to variability within turf grass species and application techniques, neither the manufacturer nor the seller has determined whether or not **INSIGNIA SC Fungicide** can safely be used on all turf grasses under all conditions. Therefore, it is recommended that the user determine if **INSIGNIA SC Fungicide** can be used safely before broad use. Apply the recommended use rate of **INSIGNIA SC Fungicide** on a small test area under conditions expected to be encountered. Monitor for any adverse effects during a 14-day period after application.

TANK MIXING

This product may be tank mixed with a fertilizer, a supplement or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact BASF Canada Inc. at 1-877-371-2273 for information before mixing any pesticide that is not specifically recommended on this label.

Physical incompatibility, reduced disease control or turf injury may result from mixing **INSIGNIA SC Fungicide** with other products. Therefore, before using any tank mix (fungicides, insecticides, liquid fertilizers, biological control products, adjuvant and additives) test a small portion of the turf to be treated to ensure that a phytotoxic response will not occur.

Additives

Do not use with organo-silicate based adjuvants or turf grass injury may occur. Due to the large number of additives or adjuvants that may be used, neither the manufacturer nor the seller has determined whether **INSIGNIA SC Fungicide** can be used safely with all additives.

Table 1. Application Rates and Intervals for Turf Grass

Disease (Pathogen)	Use Rate (mL/ 1000 sq. m.)	Use Rate (mL/ha)	Application Interval (days)	Comments
Anthracnose (<i>Colletotrichum graminicola</i>)	123 – 224	1230 – 2240	14 – 28	Apply when conditions are favourable for disease development.
Brown patch (<i>Rhizoctonia solani, zeae</i>)	112 – 200	1120 – 2000	14 – 21	Apply when conditions are favourable for disease development.
Dollar spot (<i>Sclerotinia homeocarpa</i>) - Suppression only	123 – 224	1230 – 2240	14-28	INSIGNIA SC Fungicide applied at 123 - 224 mL/1000 sq. m. for 14 to 28 day intervals will provide suppression of dollar spot.
Gray leaf spot (<i>Pyricularia grisea</i>)	112	1120	14	Apply preventative applications and continue applications while conditions are favourable for disease development.
	224	2240	28	
Gray snow Mould/Typhula blight (<i>Typhula</i> spp.)	136	1360	single application	Make a single application of 136 mL or two applications at 68 mL, 14-28 days apart in late fall just prior to snow cover.
	68	680	14 – 28	
Leaf spot (<i>Drechslera</i> spp., <i>Bipolaris</i> spp.)	123	1230	14	Apply when conditions are favourable for disease development.
Microdochium patch (Fusarium patch) and Pink snow mould (<i>Microdochium nivale</i>)	200	2000	single application	Make a single application in the late fall just prior to snow cover.
				Tankmix with PREMIS 200 F Fungicide for improved control if severe disease pressure is anticipated.
Pythium blight (<i>Pythium aphanidermatum</i>)	180	1800	14	Begin preventative application when conditions are favourable for disease development.
Pythium root dysfunction (<i>Pythium volutum, Pythium arrhenomanes, Pythium aristosporum</i>)	224	2240	14 – 28	Apply preventively for control. Irrigate immediately following application. After each application of INSIGNIA SC Fungicide rotate with at least one application of a non-group 11 fungicide registered for this disease prior to additional INSIGNIA SC Fungicide application.
Rust (<i>Puccinia</i> spp.)	100	1000	14	Apply when conditions are favourable for disease development.
Take-all patch (<i>Gaeumannomyces graminis</i>) – Suppression only	224	2240	28	Make 1 application in the fall and 1 application in the spring.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

Mode of Action

Pyraclostrobin, the active ingredient of **INSIGNIA SC Fungicide**, belongs to the group of cellular respiration inhibitors classified by the EPA and PMRA as Quinone Outside Inhibitors (QoI), or Target Site of Action Group 11 Fungicides.

Resistance Management

For resistance management, **INSIGNIA SC Fungicide** contains a Group 11 fungicide. Any (fungal/bacterial) population may contain individuals naturally resistant to **INSIGNIA SC Fungicide** and other Group 11 fungicide. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of **INSIGNIA SC Fungicide** or other Group 11 fungicides with different groups that control the same pathogens.
- Use tank mixtures with fungicide from a different group that is effective on the target pathogen when such use is permitted.
- Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- To maintain the performance of **INSIGNIA SC Fungicide** in the field, do not exceed the total number of sequential applications of **INSIGNIA SC Fungicide** and the total number of applications of **INSIGNIA SC Fungicide** per season as stated below. Do not exceed (6) six applications per season. Do not make sequential applications of **INSIGNIA SC Fungicide** or another Group 11 fungicide for control of Pythium root dysfunction, Pythium blight, Gray leaf spot or Anthracnose. Do not make more than two (2) consecutive applications of **INSIGNIA SC Fungicide** or another Group 11 fungicide for all other diseases. Alternate to an effective non-strobilurin fungicide with a different mode of action for at least two (2) applications.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Monitor treated fungal populations for resistance development. Notify BASF Canada Inc. if reduced sensitivity of the pathogen to **INSIGNIA SC Fungicide** is suspected. If disease continues to progress after treatment with this product, do not increase the use rate. Discontinue use of this product, and switch to another fungicide with a different site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.

For further information and to report suspected resistance, contact BASF Canada Inc. at 1-877-371-2273 or at www.betterturf.ca.

MIXING

1. Clean spray tank following sprayer clean-up recommendations on the previous product label.
2. Fill the spray tank one-half full of water and start agitation.
3. Add the required amount of **INSIGNIA SC Fungicide** to the tank.
4. If required, add the required amount of **PREMIS 200 F Fungicide** to tank.
5. Continue agitation while filling the remainder of the spray tank.
6. After use, clean the spray tank.

RESTRICTIONS AND LIMITATIONS

1. Do not re-enter treated area until sprays have dried.
2. Do not apply more than a six applications per season.

PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. Fatal or poisonous if swallowed. May irritate the skin. Avoid contact with skin.
3. Wash thoroughly after handling and before eating, drinking or smoking.
4. Wear protective equipment and clothing, including chemical-resistant gloves, coveralls over long-sleeved shirt, long pants, and shoes plus socks during mixing/loading, sprayer clean-up, repair and application.
5. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
6. Clean spray equipment thoroughly after use.
7. Do not contaminate domestic or irrigation water, lakes, streams or ponds by the cleaning of equipment or the disposal of wastes.

8. Apply only when the potential for drift to areas of human habitation or areas of human activity (excluding golf courses) such as houses, cottages, schools and recreation areas including parks, school grounds, and playing fields is minimal, taking into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15–20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

The patient should be treated symptomatically.

ENVIRONMENTAL PRECAUTIONS

Toxic to aquatic organisms, non-target terrestrial plants and small wild mammals. Observe buffer zones specified under DIRECTIONS FOR USE. To reduce runoff from treated areas into aquatic habitats avoid application to areas with a moderate to steep slope, compacted soil, or clay. Avoid application when heavy rain is forecast. Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

STORAGE

1. Store in original tightly closed container.
2. Do not ship or store near food, feed, seed and fertilizers.
3. Store in cool, dry, locked, well-ventilated area without floor drain.

4. Protect from freezing.
5. Store this product away from food or feed.

DISPOSAL

For Recyclable Containers

Do not reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for any further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

For Non-Returnable Containers

1. Triple or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial/territorial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial/territorial requirements.
5. For information on the disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

For Returnable-Refillable Containers

For disposal, this container may be returned to the point of purchase (distributor/dealer). It must be refilled by the distributor/dealer with the same product. Do not reuse this container for any other purpose.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Container

GROUP	3	FUNGICIDE
-------	----------	-----------

**PREMIS® 200 F
FUNGICIDE**

For the control or suppression of anthracnose, dollar spot, gray snow mold, pink snow mold, fusarium patch, red thread, rust and algae on established golf course turf only

SUSPENSION

COMMERCIAL

ACTIVE INGREDIENT: Triticonazole.....200 g/L

Contains 1,2 benzisothiazolin-3-one at 0.03% as a preservative

REGISTRATION NO. 28387

PEST CONTROL PRODUCTS ACT

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**

WARNING - EYE AND SKIN IRRITANT

READ THE LABEL AND ATTACHED BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 3.2 L

BASF Canada Inc.
5025 Creekbank Road
Building A, 2nd Floor
Mississauga, Ontario L4W 0B6
1-877-371-2273

PREMIS is a registered trade-mark of BASF Agro Trademarks GmbH, used under license by BASF Canada Inc.

PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. Causes eye irritation. **DO NOT** get in eyes. May irritate the skin. Avoid contact with skin.
3. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab.
4. Wash hands and exposed skin thoroughly after handling the concentrate and after each application, clean-up and repair.
5. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
6. AVOID WORKING IN SPRAY MIST.
7. **DO NOT** enter or allow entry into treated areas of the golf course until sprays have dried.
8. Apply only when the potential for drift to areas of human habitation or other areas of human activity (other than golf courses), such as parks, school grounds, and playing fields, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.
9. For use on established golf course turf.
10. **DO NOT** use beyond the course boundary.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

The patient should be treated symptomatically.

STORAGE

Store the product in the original container. Store the product in an area that is out of reach of children and animals, away from fertilizers and seed. Store this product away from food or feed.

DISPOSAL

For recyclable containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

For non-recyclable, non-returnable or non-re-fillable containers

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial/territorial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial/territorial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.

Booklet

GROUP	3	FUNGICIDE
-------	----------	-----------

**PREMIS® 200 F
FUNGICIDE**

For the control or suppression of anthracnose, dollar spot, gray snow mold, pink snow mold, fusarium patch, red thread, rust and algae on established golf course turf only

SUSPENSION

COMMERCIAL

ACTIVE INGREDIENT: Triticonazole.....200 g/L

Contains 1,2 benzisothiazolin-3-one at 0.03% as a preservative

REGISTRATION NO. 28387

PEST CONTROL PRODUCTS ACT

**IN CASE OF EMERGENCY ENDANGERING LIFE OR PROPERTY
INVOLVING THIS PRODUCT, CALL DAY OR NIGHT
1-800-454-2673**

WARNING - EYE AND SKIN IRRITANT

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

NET CONTENTS: 3.2 L

BASF Canada Inc.
5025 Creebank Road
Building A, 2nd Floor
Mississauga, Ontario L4W 0B6
1-877-371-2273

PREMIS is a registered trade-mark of BASF Agro Trademarks GmbH, used under license by BASF Canada Inc.

GENERAL DIRECTIONS FOR USE

APPLICATION ON ESTABLISHED GOLF COURSE TURF: **PREMIS 200 F Fungicide** is formulated for use on established golf course turf. **PREMIS 200 F Fungicide** is effective as a preventative spray. Ensure that the area to be treated is covered uniformly. Apply the required amount of **PREMIS 200 F Fungicide** in 6 to 8.0 litres of water per 100 m². When treating golf greens, always treat aprons and fairway approaches.

As this product is not registered for the control of pests in aquatic systems, **DO NOT** use to control aquatic pests.

DO NOT contaminate irrigation or drinking water supplies or aquatic habitats by cleaning of equipment or disposal of wastes.

Field sprayer application: **DO NOT** apply during periods of dead calm. Avoid application of this product when winds are gusty. **DO NOT** apply with spray droplets smaller than the American Society of Agricultural Engineers (ASAE S572.1) medium classification. Boom height must be 60 cm or less above the crop or ground.

DO NOT apply using aerial application equipment.

When applied as directed, **PREMIS 200 F Fungicide** will control the listed diseases, unless otherwise indicated as suppression.

Diseases Controlled	Use Rate (mL/100 sq. m.)	Comments
Gray Snow Mold (<i>Typhula incarnata</i> , <i>T. ishikariensis</i>) (Suppression)	16 – 21	Apply PREMIS 200 F Fungicide in late fall or early winter before snow fall. Use the higher rate if disease pressure is expected to be high.
Fusarium Patch and Pink Snow Mold (<i>Microdochium nivale</i>) (Suppression)	16 – 21	Apply PREMIS 200 F Fungicide in late fall or early winter before snow fall. Use the higher rate if disease pressure is expected to be high.
Anthraco (<i>Colletotrichum graminicola</i>)	16 – 21	Apply PREMIS 200 F Fungicide at the first sign of anthracnose and/or when environmental conditions are favourable for disease development. Use the higher rate when anthracnose pressure is high.
Dollar Spot (<i>Sclerotinia homeocarpa</i>)	16 – 21	Apply PREMIS 200 F Fungicide at the first sign of dollar spot or when environmental conditions are favourable for dollar spot development. Use the higher rate if disease pressure is high.
Red thread (<i>Laetisaria fuciformis</i>)	16 – 21	Apply when conditions are favourable for disease development when night temperatures are 15.6°C to 21.1°C and in periods of higher rainfall. Under preventive applications where light disease pressure is anticipated , use the lower rate and longer interval. Under severe disease conditions , use the higher rate.

Diseases Controlled	Use Rate (mL/100 sq. m.)	Comments
Rust (<i>Puccinia</i> spp.)	16 – 21	Apply when conditions are favourable for disease development. Under preventive applications where light disease pressure is anticipated, use the lower rate and longer interval. Under severe disease conditions, use the higher rate.
Algae (Suppression)	16 – 21	For algae suppression, use the lower rate for preventive applications. Under severe pest pressure or adverse weather that might be conducive to algae growth, use the higher rate.

Use **PREMIS 200 F Fungicide** in a rotational spray program with other different mode of action fungicides to reduce the possibility of disease resistance.

Maximum of one application per season.

TANK MIXING

This product may be tank mixed with a fertilizer, a supplement or with registered pest control products, whose labels also allow tank mixing, provided the entirety of both labels, including Directions For Use, Precautions, Restrictions, Environmental Precautions, and Spray Buffer Zones are followed for each product. In cases where these requirements differ between the tank mix partner labels, the most restrictive label must be followed. Do not tank mix products containing the same active ingredient unless specifically listed on this label.

In some cases, tank mixing pest control products can result in reduced pesticide efficacy or increased host crop injury. The user should contact BASF Canada Inc. at 1-877-371-2273 for information before mixing any pesticide that is not specifically recommended on this label.

Physical incompatibility, reduced disease control or turf injury may result from mixing **PREMIS 200 F Fungicide** with other products. Therefore, before using any tank mix (fungicides, insecticides, liquid fertilizers, biological control products, adjuvant and additives) test a small portion of the turf to be treated to ensure that a phytotoxic response will not occur.

SPRAY BUFFER ZONES

Spot treatments using hand-held equipment do not require a spray buffer zone. Use of low-clearance hooded or shielded sprayers that prevent spray contact with foliage.

The spray buffer zones specified in the table below are required between the point of direct application and the closest downwind edge of sensitive terrestrial habitats (such as grasslands, forested areas, shelter belts, woodlots, hedgerows, riparian areas and shrublands), and sensitive freshwater habitats (such as lakes, rivers, sloughs, ponds, prairie potholes, creeks, marshes, streams, reservoirs, and wetlands).

Method of application	Crop	Spray Buffer Zones (metres) Required for the Protection of:		
		Freshwater Habitat of Depths:		Terrestrial Habitat
		Less than 1 m	Greater than 1 m	
Field sprayer (groundboom)	Established golf course (turf)	3	1	4

When tank mixes are permitted, consult the labels of the tank-mix partners and observe the largest (most restrictive) spray buffer zone of the products involved in the tank mixture and apply using the coarsest spray (ASAE) category indicated on the labels for those tank mix partners.

The spray buffer zones for this product can be modified based on weather conditions and spray equipment configuration by accessing the Spray Buffer Zone Calculator on the Pesticides portion of the Canada.ca website.

ENVIRONMENTAL PRECAUTIONS

Toxic to birds and non-target terrestrial plants. Observe spray buffer zones specified under DIRECTIONS FOR USE.

Toxic to certain beneficial arthropods (soil dwelling beneficials). Minimize spray drift to reduce harmful effects on beneficial arthropods in habitats next to the application site such as hedgerows and woodland.

Toxic to aquatic organisms. Observe spray buffer zones specified under DIRECTIONS FOR USE.

To reduce runoff from treated areas into aquatic habitats, avoid application to areas with a moderate to steep slope, compacted soil, or clay.

Avoid application when heavy rain is forecast.

Contamination of aquatic areas as a result of runoff may be reduced by including a vegetative strip between the treated area and the edge of the water body.

This product demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this product in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, please note that **PREMIS 200 F Fungicide** contains a Group 3 fungicide. Any fungal population may contain individuals naturally resistant to **PREMIS 200 F Fungicide** and other Group 3 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields. Other resistance mechanisms that are not linked to site of action but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance:

- Where possible, rotate the use of **PREMIS 200 F Fungicide** or other Group 3 fungicides with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that is effective on the target pathogen when such use is permitted.
- Fungicide use should be based on an integrated disease management program that includes scouting, historical information related to pesticide use and crop rotation and considers host plant resistance, impact of environmental conditions on disease development, disease threshold, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications.
- Monitor treated fungal populations for resistance development. Notify BASF Canada Inc. if reduced sensitivity of the pathogen to **PREMIS 200 F Fungicide** is suspected.
- If disease continues to progress after treatment with this product, DO NOT increase the use rate. Discontinue use of this product, and switch to another fungicide with a different site of action, if available.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information and to report suspected resistance, contact BASF Canada Inc. at 1-877-371-2273 or at www.betterturf.ca.

MIXING

1. Ensure the spray tank is clean before use. Follow the clean-out recommendations stated on the label of the product that was previously used.
2. Fill the spray tank one-half full of water.
3. Add **PREMIS 200 F Fungicide** and agitate well.

4. Add the required amount of the tank mix partner.
5. Add the remainder of the water while continuing to agitate.
6. Ensure mixing is complete and agitate thoroughly before each application and continue to agitate during spraying. A prepared spray mixture should be used as soon as possible in the same day.
7. After use, clean the spray tank according to PRECAUTIONS.

RESTRICTIONS AND LIMITATIONS

1. **DO NOT** store **PREMIS 200 F Fungicide** near foodstuffs.
2. Thoroughly clean application equipment after each use.
3. **DO NOT** contaminate irrigation and drinking water supplies, and aquatic habitat by cleaning of equipment or disposal of wastes.
4. **DO NOT** apply beyond the course boundary.
5. **DO NOT** apply during periods of dead calm or when winds are gusty.
6. **DO NOT** apply more than 1 application per year.

PRECAUTIONS

1. **KEEP OUT OF REACH OF CHILDREN.**
2. Causes eye irritation. **DO NOT** get in eyes. May irritate the skin. Avoid contact with skin.
3. Wear a long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes during mixing, loading, application, clean-up and repair. Gloves are not required during application within a closed cab.
4. Wash hands and exposed skin thoroughly after handling the concentrate and after each application, clean-up and repair.
5. If clothing becomes contaminated, remove and wash separately from household laundry before reuse.
6. **AVOID WORKING IN SPRAY MIST.**
7. **DO NOT** enter or allow entry into treated areas of the golf course until sprays have dried.

8. Apply only when the potential for drift to areas of human habitation or other areas of human activity (other than golf courses), such as parks, school grounds, and playing fields, is minimal. Take into consideration wind speed, wind direction, temperature inversions, application equipment and sprayer settings.
9. For use on established golf course turf.
10. **DO NOT** use beyond the course boundary.

FIRST AID

If swallowed: Call a poison control centre or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control centre or doctor. Do not give anything by mouth to an unconscious person.

If on skin or clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control centre or doctor for treatment advice.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control centre or doctor for further treatment advice.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control centre or doctor for treatment advice.

Take container, label or product name and Pest Control Product Registration Number with you when seeking medical attention.

TOXICOLOGICAL INFORMATION

The patient should be treated symptomatically.

STORAGE

Store the product in the original container. Store the product in an area that is out of reach of children and animals, away from fertilizers and seed. Store this product away from food or feed.

DISPOSAL

For recyclable containers

DO NOT reuse this container for any purpose. This is a recyclable container, and is to be disposed of at a container collection site. Contact your local distributor/dealer or municipality for the location of the nearest collection site. Before taking the container to the collection site:

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Make the empty, rinsed container unsuitable for further use.

If there is no container collection site in your area, dispose of the container in accordance with provincial/territorial requirements.

For non-recyclable, non-returnable or non-re-fillable containers

1. Triple- or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial/territorial instruction for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial/territorial requirements.

For information on disposal of unused, unwanted product, contact the manufacturer or the provincial/territorial regulatory agency. Contact the manufacturer and the provincial/territorial regulatory agency in case of a spill, and for clean-up of spills.

NOTICE TO USER

This pest control product is to be used only in accordance with the directions on the label. It is an offence under the *Pest Control Products Act* to use this product in a way that is inconsistent with the directions on the label.